arXiv:math/0504097v3 [math.GR] 1 Sep 2005

# ON NORMAL SUBGROUPS OF PRODUCT OF GROUPS

ASHISH KUMAR DAS

Abstract. The object of this paper is to find a necessary and sufficient condition for the groups $G_1, G_2, \ldots, G_n$ so that every normal subgroup of the product $\prod_{i=1}^{n} G_i$ is of the type $\prod_{i=1}^{n} N_i$ with $N_i \trianglelefteq G_i$, $i = 1, 2, \ldots, n$. As a consequence we obtain a well-known result due to R. Remak about centreless completely reducible groups having finitely many direct factors.

## 1. Introduction

Let $G$ be a finite group. Then $G$ has a composition series given by $\{e\} = H_0 \lhd H_1 \lhd \cdots \lhd H_{n-1} \lhd H_n = G$. The set-with-multiplicities of composition factors associated to this composition series is given by $C(G) = \{H_i/H_{i-1} : i = 1, 2, \ldots, n\}$; noting that any two composition series of $G$ give rise to the same set-with-multiplicities of composition factors, upto isomorphism of factors (see [2], Jordan-Holder Theorem). It is a standard fact (see [1]) that $\forall\ K \lhd G$, $C(G)$ is the disjoint union (i.e. union counting multiplicities) of $C(G/K)$ and $C(K)$. By convention $C(\{e\}) = \emptyset$.

In [3], Leinster has proved that if $G_1$and $G_2$ are any two finite groups such that $C(G_1)$ and $C(G_2)$ have no member in common then every normal subgroup of $G_1 \times G_2$ is of the type $N_1 \times N_2$ where $N_i \trianglelefteq G_i$, $i = 1, 2$. However, considering $G_1 = G_2 =$ the alternating group $A_5$ (a simple non-abelian group), one can see that the converse is not true.

In this paper we derive a condition which is necessary as well as sufficient for any pair of groups $G_1$ and $G_2$ (finite or infinite, abelian or non-abelian) so that all normal subgroups of $G_1 \times G_2$ are of the above mentioned type. We then extend our result to any finite number of groups and obtain, as a consequence, a well-known result of R. Remak about centreless completely reducible groups having finitely many direct factors. The motivation for this work lies in finding examples of non-abelian perfect groups; a perfect group being one in which the

---

1991 *Mathematics Subject Classification.* 20D06, 20D40.

*Key words and phrases.* normal subgroups, composition series, simple groups.

sum of the orders of all normal subgroups equals twice the order of the group (see [3]).

## 2. The NS-condition

We begin this section with a few definitions.

**Definition 2.1.** Let $G_1$ and $G_2$ be any two groups. A normal subgroup $N$ of $G_1 \times G_2$ is said to be of standard type if $N = N_1 \times N_2$ where $N_i \trianglelefteq G_i$, $i = 1, 2$.

**Definition 2.2.** Two groups $G_1$ and $G_2$ are said to have a subgroup in common if there exist non-trivial subgroups $H_i$ of $G_i$, $i = 1, 2$, such that $H_1 \cong H_2$.

**Definition 2.3.** Two groups $G_1$ and $G_2$ are said to satisfy NS-condition if $\forall\ H_i \triangleleft G_i$, $i = 1, 2$, the centres $Z(G_1/H_1)$ and $Z(G_2/H_2)$ of the quotient groups $G_1/H_1$ and $G_2/H_2$ have no subgroup in common.

**Remark 2.4.** If $H \trianglelefteq G$ then the subgroups of $Z(G/H)$ are of the type $K/H$ where $H \trianglelefteq K \leq G$ and $gkg^{-1}k^{-1} \in H, \quad \forall g \in G$ and $\forall k \in K$. In particular $K \trianglelefteq G$.

The following proposition shows that NS-condition is strictly weaker than Leinster's hypothesis (see Proposition 3.1 of [3]).

**Proposition 2.5.** *If $G_1$ and $G_2$ are two finite groups such that $C(G_1)$ and $C(G_2)$ have no abelian member in common then $G_1$ and $G_2$ satisfy NS-condition. However, the converse is not true.*

*Proof.* Suppose $H_i \triangleleft G_i$, $i = 1, 2$, are such that the centres $Z(G_1/H_1)$ and $Z(G_2/H_2)$ have a subgroup in common. So, there are non-trivial subgroups $K_i/H_i$ of $Z(G_i/H_i)$, $i = 1, 2$, such that $K_1/H_1 \cong K_2/H_2$. Then $C(K_1/H_1) = C(K_2/H_2)$. Since $H_i \triangleleft K_i \trianglelefteq G_i$, we have $C(K_i/H_i) \subseteq C(K_i) \subseteq C(G_i)$, $i = 1, 2$. Thus, $C(G_1)$ and $C(G_2)$ have an abelian member in common; noting that both $K_1/H_1$ and $K_2/H_2$ are abelian.

That the converse is not true can be easily seen by considering $G_1 = S_4$ and $G_2 = \mathbb{Z}/3\mathbb{Z}$, and noting that $S_4$ and $S_4/V \cong S_3$ have trivial centres; $V$ being the normal subgroup $\{e, (12)(34), (13)(24), (14)(23)\}$ of $S_4$. □

**Proposition 2.6.** *Two finite groups $G_1$ and $G_2$ satisfy NS-condition if and only if $gcd(n_1, n_2) = 1$ where*

$$n_i = \prod_{H_i \trianglelefteq G_i} |Z(G_i/H_i)|, \quad i = 1, 2.$$

*Proof.* If each $n_i$ is divisible by a prime $p$, then p divides $|Z(G_i/H_i)|$ for some $H_i \trianglelefteq G_i$, $i = 1, 2$. So, each of $Z(G_i/H_i)$, $i = 1, 2$, has a subgroup of order $p$ (see [1]) . Obviously such a subgroup is isomorphic to $\mathbb{Z}/p\mathbb{Z}$.

Conversely, if $gcd(n_1, n_2) = 1$ then $\forall\ H_i \trianglelefteq G_i$, $i = 1, 2$, the centres $Z(G_1/H_1)$ and $Z(G_2/H_2)$ have coprime orders and so they can not have any subgroup in common. □

The following corollary is immediate.

**Corollary 2.7.** *If two finite groups $G_1$ and $G_2$ have coprime orders then they satisfy NS-condion. Converse also holds if $G_1$ and $G_2$ are finite abelian.*

## 3. Main Theorem

In this section we show that the NS-condition is indeed necessary as well as sufficient.

**Theorem 3.1.** *Let $G_1$ and $G_2$ be any two groups (finite or infinite, abelian or non-abelian). Then all normal subgroups of $G_1 \times G_2$ are of standard type if and only if $G_1$ and $G_2$ satisfy NS-condition.*

*Proof.* Suppose $G_1$ and $G_2$ satisfy NS-condition. Let $N \trianglelefteq G_1 \times G_2$. Set $H_1 = \pi_1((G_1 \times \{e_2\}) \cap N)$ and $H_2 = \pi_2((\{e_1\} \times G_2) \cap N)$ where $e_i$ are identities of $G_i$ and $\pi_i : G_1 \times G_2 \longrightarrow G_i$ are projections, $i = 1, 2$. Then $H_1 \times \{e_2\}, \{e_1\} \times H_2 \subset N \subset \pi_1(N) \times \pi_2(N)$ and so

$$H_1 \times H_2 = (H_1 \times \{e_2\})(\{e_1\} \times H_2) \subset \pi_1(N) \times \pi_2(N) \tag{3.2}$$

It may be noted here that $H_i \trianglelefteq G_i$ and $H_i \trianglelefteq \pi_i(N)$, $i = 1, 2$. Now, suppose $a_1 \in \pi_1(N)$. Then $(a_1, a_2) \in N$ for some $a_2 \in G_2$; in fact $a_2 \in \pi_2(N)$. Therefore, $\forall g_1 \in G_1$, we have

$$\begin{aligned} &(g_1 a_1 {g_1}^{-1}, a_2) = (g_1, e_2)(a_1, a_2)({g_1}^{-1}, e_2) \in N \\ \Longrightarrow\ &(g_1 a_1 {g_1}^{-1} {a_1}^{-1}, e_2) \in N \\ \Longrightarrow\ &g_1 a_1 {g_1}^{-1} {a_1}^{-1} \in H_1 \\ \Longrightarrow\ &g_1 H_1 a_1 H_1 = a_1 H_1 g_1 H_1 \in G_1. \end{aligned}$$

Thus, $a_1 H_1 \in Z(G_1/H_1)$. So, we have $\pi_1(N)/H_1 \subset Z(G_1/H_1)$. Similarly, $\pi_2(N)/H_2 \subset Z(G_2/H_2)$. Note that if $a_1, b_1 \in \pi_1(N)$ then $(a_1, a_2), (b_1, b_2) \in N$ for some $a_2, b_2 \in \pi_2(N)$, and so $(a_1 {b_1}^{-1}, a_2 {b_2}^{-1})$, $(a_1 b_1, a_2 b_2) \in N$. Therefore,

$$\begin{aligned} a_1 H_1 = b_1 H_1 \iff a_1 {b_1}^{-1} \in H_1 \iff (a_1 {b_1}^{-1}, e_2) \in N \\ \iff (e_1, a_2 {b_2}^{-1}) \in N \iff a_2 {b_2}^{-1} \in H_2 \iff a_2 H_2 = b_2 H_2. \end{aligned}$$

This means that we have a well-defined injective map $f : \pi_1(N)/H_1 \longrightarrow \pi_2(N)/H_2$ given by $f(a_1H_1) = a_2H_2$ where $(a_1, a_2) \in N$. Also, $f(a_1H_1b_1H_1) = f(a_1b_1H_1) = a_2b_2H_2 = a_2H_2b_2H_2 = f(a_1H_1)f(b_1H_1)$, showing that $f$ is a homomorphism. Finally, if $b \in \pi_2(N)$ then $(a, b) \in N$ for some $a \in \pi_1(N)$ and so $f(aH_1) = bH_2$, which implies that $f$ is surjective. Thus $f$ is an isomorphism. Hence it follows from the hypothesis that $\pi_i(N)/H_i$ are trivial subgroups of $Z(G_i/H_i)$, $i = 1, 2$. Therefore, $H_i = \pi_i(N)$, $i = 1, 2$, and so $N = H_1 \times H_2$, by (3.2).

Conversely, suppose $G_1$ and $G_2$ do not satisfy the NS-condition. So, there exist $H_i \lhd G_i$, $i = 1, 2$, such that $Z(G_1/H_1)$ and $Z(G_2/H_2)$ have a subgroup in common. Let $K_i/H_i$ be non-trivial subgroups of $Z(G_i/H_i)$, $i = 1, 2$, such that there is an isomorphism $F : K_1/H_1 \longrightarrow K_2/H_2$. Put $N = \{(a_1, a_2) \in K_1 \times K_2 : F(a_1H_1) = a_2H_2\}$. Let $(a_1, a_2), (b_1, b_2) \in N$ then $F(a_1H_1) = a_2H_2$ and $F(b_1H_1) = b_2H_2$. So, $F(a_1{b_1}^{-1}H_1) = a_2{b_2}^{-1}H_2$. Thus $(a_1, a_2)(b_1, b_2)^{-1} = (a_1{b_1}^{-1}, a_2{b_2}^{-1}) \in N$, showing that $N$ is a subgroup of $G_1 \times G_2$. Again let $(a_1, a_2) \in N$ and $(g_1, g_2) \in G_1 \times G_2$. Then, $(g_1, g_2)(a_1, a_2)(g_1, g_2)^{-1} = (g_1a_1{g_1}^{-1}, g_2a_2{g_2}^{-1}) \in K_1 \times K_2$, since, by Remark 2.4, $K_i \trianglelefteq G_i$, $i = 1, 2$. Also, since $a_iH_i \in K_i/H_i \subset Z(G_i/H_i)$, $i = 1, 2$, we have

$$F(g_1a_1{g_1}^{-1}H_1) = F(a_1H_1) = a_2H_2 = g_2a_2{g_2}^{-1}H_2.$$

Thus $(g_1, g_2)(a_1, a_2)(g_1, g_2)^{-1} \in N$, and so $N \trianglelefteq G_1 \times G_2$. On the other hand, suppose $N$ is of standard form $N_1 \times N_2$ where $N_i \trianglelefteq G_i$, $i = 1, 2$. Then, $\pi_i(N) = N_i$, $i = 1, 2$. But since F is bijective, we have $\pi_i(N) = K_i$, $i = 1, 2$. Therefore, $N = K_1 \times K_2$. Since $K_1/H_1$ is non-trivial, there is some $a_1 \in K_1$ such that $a_1H_1 \neq H_1$. But $(a_1, e_2) \in K_1 \times K_2 = N$. So, $F(a_1H_1) = e_2H_2 = H_2$, the zero element of $K_2/H_2$. Therefore, since $F$ is injective, we have $a_1H_1 = H_1$, the zero element of $K_1/H_1$. This contradiction shows that $N$ is not of standard type. □

Note that $(G_1 \times G_2)/(H_1 \times H_2) \cong G_1/H_1 \times G_2/H_2$ where $H_i \trianglelefteq G_i$, $i = 1, 2$. Therefore, using induction and the fact that the centre of a product of groups equals the product of the centres of the constituent groups, we have the following corollary:

**Corollary 3.3.** *Let $G_1, G_2, \ldots, G_n$ be any $n$ groups (finite or infinite, abelian or non-abelian). Then every normal subgroup of the product $\prod_{i=1}^{n} G_i$ is of the type $\prod_{i=1}^{n} N_i$ with $N_i \trianglelefteq G_i$, $i = 1, 2, \ldots, n$, if and only if $G_1, G_2, \ldots, G_n$ satisfy NS-condition pairwise.*

**Remark 3.4.** If atleast $n - 1$ of the groups $G_1, G_2, \ldots, G_n$ are simple non-abelian then they satisfy NS-condition pairwise. In view of this, the above corollary says, in particular, that if G is a direct product of non-abelian simple groups $G_i$, $i = 1, 2, \ldots, n$, then every normal subgroup of $G$ is a direct product of certain of the $G_i$'s. This is a well-known result proved by R. Remak (see [4], page 88).

## References

[1] J. B. Fraleigh, *A first course in Abstract Algebra*, Addison-Wesley Publishing Company, Inc. Philippines, 1982.

[2] N. Jacobson, *Basic Algebra - vol I*, W. H. Freeman and Company, U.S.A., 1974

[3] T. Leinster,*Perfect number and groups*, arXiv:math.GR/0104012 v1 Apr 2001.

[4] Derek J. S. Robinson, *A course in the Theory of Groups* (Second Edition), Graduate Text in Mathematics 80, Springer, New York, 1996.

Department of Mathematics, North Eastern Hill University, Permanent Campus, Shillong-793022, Meghalaya, India.

*E-mail address*: `akdas@nehu.ac.in`